

ANTIHYPERTENSIVES

KEY POINTS

- Lowering blood pressure reduces risk of a range of long term consequences, this benefit is still evident in the elderly.
- More aggressive control of blood pressure in the elderly may be equivalent in benefit to less aggressive control of hypertension.
- Low blood pressure may be associated with increased morbidity and mortality in the elderly.
- Patients being treated for hypertension are more likely to fall if they have proven postural hypotension.
- Adverse effects of many antihypertensive agents are likely to be more common in the elderly.
- Withdrawal of antihypertensives should be gradual.

CONTEXT

This guide considers the deprescribing of antihypertensive medications. Caution is required when patients have multiple morbidities as some of these agents may have a dual therapeutic indication.

RECOMMENDED DEPRESCRIBING STRATEGY

- 1 Where patients are receiving multiple agents that lower blood pressure, strategies for reduction and cessation should focus on one agent at a time.

Reduction or cessation of antihypertensive agents should be considered:
 - In frail elderly and/or immobile patients
 - In patients with a high falls risk
 - In patients with confirmed postural hypotension (>20mmHg fall in systolic on standing, and/or >10mmHg fall in diastolic on standing)
- 2
- 3 It is recommended that most antihypertensives be tapered at approximately 25% per month over 3 – 4 months.

EFFICACY

Multiple studies have demonstrated that there are increased rates of morbidity and mortality in patients with hypertension. Where appropriate treatment is provided, there is a reduction in morbidity and mortality. However, the benefit of lowering blood pressure is attenuated with increasing age. In 2002, Lewington et al published data from over 1 million adults from 61 studies on the associations between SBP and DBP and mortality from stroke and coronary heart disease by age. The associations between both SBP and DBP and mortality from stroke, coronary heart disease and other vascular disease were graded and continuous with the lowest risk at SBP of 115mm Hg and DBP of 75mm Hg (lower BP levels were not reported) and the highest risk at SBP of 175mm Hg and DBP of 105mm Hg (higher levels were not reported). However, these associations were weaker at older age (see Figure 1 on page 2).¹

Trials of hypertension management in the elderly are limited and were reviewed by Fleg et al.² in 2011 and Muntner et al.³ in 2014. Of 12 studies reviewed by Fleg, five showed statistically significant reductions in cardiovascular events. The key points from these five studies were:

- Mean ages between 67-72 years, with between 40 and 58% females
- Entry BPs were from 169-185 systolic
- All five studies showed a relative risk reduction of stroke between 23 and 57%

Muntner's article reviews three other papers that targeted intensive vs more lenient systolic blood pressure control in older patients:

- A Japanese study of 4418 patients aged 65-84years compared tight vs lenient control of blood pressure on outcomes. One group achieved

continued page 2 >




Figure 1: Stroke mortality in each decade of age versus usual blood pressure³

DRUG CLASS (COMMON EXAMPLES)	ADVERSE EFFECTS
Thiazide and Loop diuretics (hydrochlorothiazide, indapamide, chlorthalidone, frusemide)	Hypokalemia, hyponatremia, hypomagnesemia Volume-depletion and orthostatic hypotension Renal impairment, hyperuricemia, gout, lipid alterations, hyperglycemia, insulin resistance NSAIDs reduce thiazide potency Erectile dysfunction and possibly impotence Reduction of lithium excretion and precipitate lithium toxicity Potential to increase fatigue and lethargy Pro-diabetogenic potential in combination with Beta Blockers Increase of urinary frequency, leg cramps Decrease of renal blood flow, creatinine clearance, Glomerular Filtration Rate
Potassium Sparing Diuretics (spironolactone, amiloride)	Hyperkalemia, hypotension
Beta Blockers (atenolol, metoprolol, propranolol)	Sinus bradycardia, fatigue, AV-nodal heart block bronchospasm, aggravation of acute heart failure Intermittent claudication, confusion, hyperglycemia Diabetes mellitus Drowsiness, lethargy, sleep disturbance, visual hallucinations, depression, blurring of vision, nightmares Pulmonary side-effects (increased airway resistance in asthmatics) Peripheral vascular side-effects (cold extremities, Raynaud's phenomenon) Erectile dysfunction
Angiotensin Converting Enzyme Inhibitors (perindopril, ramipril, fosinopril, trandolapril)	Cough, hyperkalemia Angioneurotic edema Rash, altered taste sensation, renal impairment
Angiotensin Receptor Blockers (candesartan, irbesartan)	Hyperkalemia, renal impairment
Calcium Channel Blockers (non-dihydropyridines eg. verapamil, diltiazem)	Rash, sinus bradycardia, heart block, heart failure, constipation (verapamil), gingival hyperplasia Ankle edema, headache and postural hypotension
Calcium Channel Blockers (dihydropyridines eg. amlodipine, nifedipine)	Peripheral edema, heart failure, tachycardia Aggravation of angina pectoris (short-acting agents)
Direct vasodilators (hydralazine)	Tachycardia, fluid retention Angina pectoris
Alpha 1 adrenergic blockers (prazosin)	Hypotension
Alpha-beta adrenergic blockers (labetalol)	Hypotension, heart block, sinus bradycardia, bronchospasm
Central acting agents (moxonidine, methylidopa)	Sedation, constipation, dry mouth

Table 1: Most common drug-related side effects of antihypertensive classes¹⁷

< EFFICACY continued from page 1

136/75 on average while the other 146/78. Over 2 years of follow-up, there were no differences in the primary composite outcome of cardiovascular disease or renal failure.⁴

- The Valsartan in the Elderly with Isolated Systolic Hypertension (VALISH) study found no difference in 70-84 year olds that achieved 137 vs 142 systolic BPs in terms of stroke, sudden death or myocardial infarction frequency.⁵
- An Italian study of 1111 patients with a mean age of 67 years randomised patients to tight (<130) vs moderate (<140) control of blood pressure. They showed a difference in a composite endpoint of CVD/renal disease after 2 years of 9.4% in the moderate control group vs 4.8% (ARR 4.6%, NNT= 22 over 2 years).⁶

One additional study (HYVET)) looked specifically at patients over 80 years of age.⁷ This group randomised patients with a starting systolic BP of 160mmHg or more to indapamide or a placebo. Perindopril was added to the indapamide if the target BP of 150mmHg was not achieved. They reported positive outcomes for the following endpoints after an average 1.8 year follow-up:

- **Death from stroke**
51/1933 (1.4%) vs 42/1912 (2.2%); OR 0.61 [0.38-0.99]; ARR 0.8% (NNT= 125)
- **Death from any cause**
196/1933 (10.1%) vs 235/1912 (12.3%); OR 0.79 [0.65-0.95]; ARR 2.2% (NNT= 45)
- **Development of heart failure**
22/1933 (1.14%) vs 57/1912 (2.97%); OR 0.36 [0.22-0.58]; ARR 1.83% (NNT= 54)
- **Any cardiovascular event**
138/1933 (7.14%) vs 193/1912 (10.09%); OR 0.66 [0.53-0.82]; ARR 2.95% (NNT= 34)

Thus there seem to be benefits in reducing blood pressure from high systolic blood pressures in the elderly. These relative benefits may be somewhat less than those achieved by a similar level of reduction in younger patients. However, because of increased absolute risks in the elderly, the absolute benefits may be equal to or greater than younger patients. There is also some support for the contention that more moderate blood pressure targets achieve a similar benefit to more intensive control in the elderly. It should be noted that these studies all include relatively fit elderly patients and that frail elderly patients may be more sensitive to the impact of antihypertensive treatment and may or may not obtain the same benefit from antihypertensive therapy.

In an observational study of 2340 adults older than 65 years, the association between blood pressure and mortality was examined according to whether or not individuals were frail (defined as an inability to walk 6 meters in less than 8 seconds).⁸ Among frail adults, there was no association between blood pressure and mortality. In addition, a higher blood pressure was associated with a lower risk of death among the most frail (ie, those who could not walk the distance at all). This was an observational study and there are no randomised controlled trials of treatment of hypertension in frail elderly people.

ADVERSE EFFECTS

SUSTAINED HYPOTENSION

Some studies have reported an increased cardiovascular risk at very low systolic or diastolic blood pressures in the elderly.^{9, 10, 11} Voko et al. reported a J shaped relationship between incidence of stroke and diastolic (but not systolic) blood pressure in treated hypertensives.⁹ In patients receiving treatment for hypertension, diastolic blood pressure of less than 65mmHg was associated with the same stroke risk as patients with a diastolic of >84mmHg, and significantly higher than those with a systolic of 65-74mmHg.

Ogihara et al undertook a 3 year follow-up 2164 patients over 60 who regularly attended their clinician and documented cardiovascular morbidity and mortality as well as achieved blood pressure.¹⁰ In the subgroup of patients aged 75 years or more, patients with an achieved systolic BP of <120mmHg had a significantly higher incidence of total cardiovascular events, as did patients with a systolic BP of >160mmHg (see Table 2).

Taken together, the above studies indicate that low blood pressure may result in increased morbidity and mortality in the elderly. It remains unclear whether the low blood pressure is itself an indicator of poor cardiovascular health, which may be responsible for this observation, furthermore, a reanalysis of the PROGRESS trial, that originally proposed a J curve in patients treated for hypertension concluded that lower blood pressure levels were safe and not associated with a higher stroke risk.¹²

POSTURAL HYPOTENSION

A further potential limiting factor in the treatment of elderly patients is the presence of, or exacerbation by treatment of, postural hypotension. Postural and/or postprandial hypotension is found in up to 20% of elderly patients with isolated systolic hypertension.^{13, 14} Hypertensive older patients with postural hypotension (particularly those with less well controlled hypertension) are more likely to fall than patients without.¹⁵ Indeed, antihypertensive treatment has been associated with a 43% increased risk of hip fractures in the elderly in the first 45 days of treatment.¹⁶ Measurement of lying and standing blood pressures are essential prior to commencing or modifying antihypertensive therapy.

OTHER ADVERSE EFFECTS OF ANTIHYPERTENSIVE AGENTS

There are a wide range of differing antihypertensive agents available with multiple mechanisms of action. As a result, there are a wide range of possible adverse effects from these agents, either alone or in combination with other agents being taken by the patient. Metabolic, cardiac and renal effects are seen from many of the antihypertensives, with some agents also exhibiting more specific adverse effects. For the vast majority of adverse effects, the elderly and those with limited reserve are more likely to sustain adverse effects. A summary of the complexity in choosing between drug classes for treatment of hypertension has been undertaken.¹⁷ In Table 1, the main adverse effects with each class of agents is listed.

FACTORS TO CONSIDER BEFORE DEPRESCRIBING

IN FAVOUR OF DEPRESCRIBING

- ✓ The benefits of lifestyle modification should not be underestimated. In patients where lifestyle modification (exercise, salt and sugar restriction, alcohol, weight loss) are possible, these changes can support the reduction and/or cessation of antihypertensive agents.
- ✓ The benefits of treating hypertension in the oldest old (85 and older) are unclear, ongoing treatment should be reassessed in light of prognosis, comorbidities and quality of life.
- ✓ Patients who are frail and have a high falls risk are more likely to fall as a result of antihypertensive treatment and may not derive the same benefit of treatment as non-frail elderly. Reduction or cessation of antihypertensives should be considered in these patients.

AGAINST DEPRESCRIBING

- ✗ Agents with an antihypertensive effect may have other benefits in patients with other comorbidities and they may be prescribed more specifically for these other purposes. Beta blockers for heart failure, atrial fibrillation or ischaemic heart disease, ACE inhibitors for heart failure or renal protection and prazosin for prostatic symptoms are examples of where cessation of these agents may worsen the underlying condition.

		<75 YEARS OLD		≥ 75 YEARS OLD	
BLOOD PRESSURE ACHIEVED		NUMBER OF PATIENTS	RELATIVE RISK (*=STATISTICALLY SIGNIFICANT)	NUMBER OF PATIENTS	RELATIVE RISK (*=STATISTICALLY SIGNIFICANT)
Diastolic	<70	158	1.21	98	1.77
	70-79	503	1.00 (reference)	275	1.00 (reference)
	80-89	677	0.99	240	1.02
	>89	161	1.36	52	1.76
Systolic	<120	74	0.43	34	3.40*
	120-139	740	1.00 (reference)	298	1.00 (reference)
	140-159	581	1.49	284	1.74
	>159	104	2.14	49	2.90*

Table 2: Relation between achieved blood pressure and cardiovascular events by age.¹⁰



DISCONTINUATION SYNDROMES

Withdrawal effects may be wide ranging, depending on the specific class of agent and any other conditions being treated. These may include peripheral oedema, tachycardia, rebound hypertension or worsening heart failure or ischaemic heart disease. As a result, it is recommended that most antihypertensives should be tapered at approximately 25% every month over 3-4 months.

RESOURCES

- ☒ QUICK REFERENCE GUIDE
- ☒ ANTIHYPERTENSIVES GUIDE
- ☒ ANTIPLATELET AGENTS GUIDE
- ☒ ANTIPSYCHOTICS GUIDE
- ☒ BENZODIAZEPINES GUIDE
- ☒ BISPHOSPHONATES GUIDE
- ☒ STATINS GUIDE
- ☒ VITAMIN D & CALCIUM GUIDE

REFERENCES

1. Lewington S, Clarke R, Qizilbash N, et al. Age-specific relevance of usual blood pressure to vascular mortality: a meta-analysis of individual data for one million adults in 61 prospective studies. *Lancet* 2002;360:1903-13.
2. Fleg JL, Aronow WS, Frishman WH. Cardiovascular drug therapy in the elderly: benefits and challenges. *Nat Rev Cardiol*. 2011 Jan;8(1):13-28.
3. Muntner P et al. Systolic blood pressure goals to reduce cardiovascular disease among older adults. *Am J Med Sci* 2014 48(2): 129-134.
4. JATOS Study Group. Principal results of the Japanese trial to assess optimal systolic blood pressure in elderly hypertensive patients (JATOS). *Hypertens Res* 2008;31:2115-27.
5. Ogihara T, Saruta T, Rakugi H, et al; Valsartan in Elderly Isolated Systolic Hypertension Study Group. Target blood pressure for treatment of isolated systolic hypertension in the elderly: valsartan in elderly isolated systolic hypertension study. *Hypertension* 2010;56:196-202.
6. Verdecchia P, Staessen JA, Angeli F, et al. Usual versus tight control of systolic blood pressure in non-diabetic patients with hypertension (Cardio-Sis): an open-label randomised trial. *Lancet* 2009;374:525-33.
7. The HYVET Study Group. Treatment of Hypertension in Patients 80 Years of Age or Older. *N Engl J Med* 2008;358:1887-98.
8. Odden MC, Peralta CA, Haan MN, Covinsky KE. Rethinking the association of high blood pressure with mortality in elderly adults: the impact of frailty. *Arch Intern Med*. 2012;172(15):1162.
9. Voko Z, Bots ML, Hofman A, et al. J-shaped relation between blood pressure and stroke in treated hypertensives. *Hypertension* 1999;34: 1181-5.
10. Ogihara T, Matsuoka H, Rakugi H. Practitioner's trial on the efficacy of antihypertensive treatment in elderly patients with hypertension II (PATE-hypertension II study) in Japan. *Geriatr Gerontol Int* 2011;11: 414-21.
11. Shiva Satish et al. The Relationship Between Blood Pressure and Mortality in the Oldest Old. *J Am Geriatr Soc* 2001 49: 367-74.
12. Hisatomi Arimaa, John Chalmers, Mark Woodward, Craig Anderson, Anthony Rodgers, Stephen Davis, Stephen McMahon, Bruce Neale for the PROGRESS Collaborative Group. Lower target blood pressures are safe and effective for the prevention of recurrent stroke: the PROGRESS trial. *Journal of Hypertension* 2006, 24:1201-1208.
13. Vanhanen H, Thijs L, Birkenhäger W, Tilvis R, Sarti C, Tuomilehto J, Bulpitt C, Fagard R, Staessen JA. Associations of orthostatic blood pressure fall in older patients with isolated systolic hypertension. *Syst-Eur Investigators. J Hypertens*. 1996;14(8):943.
14. Applegate WB, Davis BR, Black HR, Smith WM, Miller ST, Burlando AJ. Prevalence of postural hypotension at baseline in the Systolic Hypertension in the Elderly Program (SHEP) cohort. *J Am Geriatr Soc*. 1991;39(11):1057.
15. Gangavati A, Hajjar L, Quach L, Jones RN, Kiely DK, Gagnon P, Lipsitz LA. Hypertension, orthostatic hypotension, and the risk of falls in a community-dwelling elderly population: the maintenance of balance, independent living, intellect, and zest in the elderly of Boston study. *J Am Geriatr Soc*. 2011 Mar;59(3):383-9.
16. Butt DA, Mamdani M, Austin PC, Tu K, Gomes T, Glazier RH. The risk of hip fracture after initiating antihypertensive drugs in the elderly. *Arch Intern Med*. 2012;172(22):1739.
17. Kaiser EA, Lotze U, Schäfer HH. Increasing complexity: which drug class to choose for treatment of hypertension in the elderly? *Clin Interv Aging*. 2014 Mar 24;9:459-75.



www.consultantpharmacyservices.com.au



www.primaryhealthtas.com.au